



Documents

Ahmad, B.^a, Khan, R.A.^b, Sivasundaram, S.^c

K-th order monotone iterative scheme for boundary value problems associated with second order nonlinear functional differential equations

(2005) *Communications in Applied Analysis*, 9 (3-4), pp. 317-325. Cited 2 times.

^a Department of Mathematics, Faculty of Science, King Abdul Aziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia

^b Department of Mathematics, Quaid-i-Azam University, Islamabad, Pakistan

^c Department of Mathematics, Embry-Riddle Aeronautical University, Daytona Beach, FL 32114, United States

Abstract

The method of quasilinearization, coupled with the method of upper and lower solutions, is applied to a nonlinear second order boundary value problem associated with a functional differential equation to obtain a sequence of approximate solutions converging monotonically and rapidly to a unique solution of the problem. © Dynamic Publishers, Inc.

Document Type: Article

Source: Scopus

About Scopus

[What is Scopus](#)
[Content coverage](#)
[What do users think](#)
[Latest](#)
[Tutorials](#)

Contact and Support

[Contact and support](#)
[Live Chat](#)

About Elsevier

[About Elsevier](#)
[About SciVerse](#)
[About SciVal](#)
[Terms and Conditions](#)
[Privacy Policy](#)



Copyright © 2012 Elsevier B.V. All rights reserved. SciVerse® is a registered trademark of Elsevier Properties S.A., used under license. Scopus® is a registered trademark of Elsevier B.V.